

Simulation of SU(2) Dynamical Fermion at Finite Chemical Potential and at Finite Temperature^a

Y.Liu¹, O.Miyamura¹, A.Nakamura¹, T.Takaishi²

¹*Hiroshima University, Higashi-Hiroshima 739-8521, Japan*

²*Hiroshima University of Economics, Hiroshima 731-01 Japan*

SU(2) lattice gauge theory with dynamical fermion at non-zero chemical potential and at finite temperature is studied. We focus on the influence of chemical potential for quark condensate and mass of pseudoscalar meson at finite temperature. Hybrid Monte Carlo simulations with $N_f = 8$ staggered fermions are carried out on $12 \times 12 \times 24 \times 4$ lattice. At $\beta = 1.1$ and $m_q = 0.05, 0.07, 0.1$, we calculate the quark condensate and masses of pseudoscalar meson consisting of light and heavier quarks for chemical potential $\mu = 0.0, 0.02, 0.05, 0.1, 0.2$.

1 INTRODUCTION

The study of strong interactions at finite baryon density has a long history. Recent theoretical speculation is that the ground state of dense baryonic matter may be more exotic^{1, 2}. Nambu-Jona-Lasinio type and instanton liquid models^{2, 3} suggest that the QCD vacuum at sufficiently high baryon density could become a color superconductor³.

Considerable progress has been made in simulating lattice quantum chromodynamics (LQCD) at nonzero temperature⁴. But, there has been little progress in understanding the theory at nonzero chemical potential even though a pioneering work was presented sixteen years ago⁵. The basic difficulty in simulation of LQCD at finite density is that the effective action becomes complex due to the introduction of chemical potential. Standard algorithms such as hybrid molecular dynamics (HMD)⁶ or hybrid Monte Carlo (HMC)⁷ can not be applied in this situation. In order to avoid the complex determinant, early simulations used quenched approximation, but they do not correspond to the real world⁸. In this situation, a possible way is to examine response of physical quantities to chemical potential at $\mu = 0$. Quark-number susceptibilities have been studied by S.Gottlieb *et al*⁹ and give us the signals for the chiral- symmetry-restoration phase transition. Study of response of hadron masses to chemical potential is now under investigation by *QCD – TARO* collaboration¹⁰.

Another way is to study LQCD for color SU(2) system where action remains real. Hamiltonian approach at zero temperature and finite density has

^aTalk presented by Y.Liu

E-mail: liu@hirohe.hepl.hiroshima-u.ac.jp

been studied by E.B.Gregory *et al*¹¹, and gives us some results for vacuum energy, chiral condensate, baryon-number density and its susceptibility. Numerical simulation at non zero chemical potential has been performed by S.Hands *et al*¹², by using several algorithms; HMD algorithm ; HMC algorithm ; Two-Step Multi-Boson (TSMB) algorithm¹³ on rather small lattices (mostly on 6^4 lattice) at zero temperature. They studied relation of chiral condensate, pion mass and chemical potential at $\mu > \frac{m_\pi}{2}$ as well as $\mu < \frac{m_\pi}{2}$.

In this paper, we focus on the effects of chemical potential to masses and condensate at finite temperature ($\mu=0.02, 0.05, 0.1$ and 0.2) on larger lattices $12^3 \times 4$ and $12^2 \times 24 \times 4$. One of interests is precise study of masses of meson with nonzero chemical potential since QCD Sum Rule suggests linear dependence on density.¹⁴ Firstly, we shall observe chiral condensate with nonzero chemical potential near critical temperature. Secondly, we shall examine meson masses consisting of light-light quarks and light-heavy quarks. The organization of the paper is as follows: In Section 2 we state our procedure. Simulation is described in section 3. We show some preliminary results of the present simulation in Section 4. Summary and conclusions are stated in Section 5.

2 STAGGERED FERMION ACTION AT NONZERO CHEMICAL POTENTIAL

At non zero chemical potential, staggered fermion matrix M is given by

$$M_{x,y}(U,\mu) = 2m\delta_{xy} + \sum_{\nu=1}^3 \eta_\nu(x)[U_{x,\nu}\delta_{y,x+\hat{\nu}} - U_{y,\nu}^\dagger\delta_{y,x-\hat{\nu}}] + \eta_4(x)[e^\mu U_{x,4}\delta_{y,x+\hat{4}} - e^{-\mu}U_{y,4}^\dagger\delta_{y,x-\hat{4}}], \quad (1)$$

where U is SU(2) link variables and μ is chemical potential. η_ν is staggered phase and m is quark mass parameter.

The matrix M satisfies the following relation

$$M_{x,y}(U,\mu) = (-1)^{sig(x)-sig(y)} M_{x,y}^\dagger(U,-\mu). \quad (2)$$

The path integral for the partition function is given by

$$Z = \int [DU] \det(M)^{N_f/4} e^{-S_g}, \quad (3)$$

where S_g is plaquette action written as ,

$$S_g = \beta \sum_P [1 - \frac{1}{4} Tr(U_P + U_P^\dagger)], \quad (4)$$

where $\beta = \frac{4}{g^2}$.

In cases of simulations by HMC algorithm, we need positive definiteness of hamiltonian. Since $\det(M)$ is real for SU(2) system, the positive definiteness condition is achieved by taking $\det(M)^2$ which includes $n_f = 8$ dynamical fermions. Note that the standard even-odd partitioning of the fermion matrix which is commonly used to reduce the number of flavors can not be applied for simulations at finite chemical potential. Thus $n_f = 8$ is the minimum number of flavors for SU(2) HMC simulations with dynamical fermions. The present study is performed with $n_f = 8$.

3 SIMULATIONS

Firstly, we have done simulation on 6^4 lattice and confirmed that our results are consistent with the previous work by S.Hands *et al*¹². After that, numerical simulations on $12^3 \times 4$ and $12^2 \times 24 \times 4$ lattices have been carried out for parameters summarized in Table 1.

Table 1: List of the data sample

N_F	m_q	μ	lattice size	β
8	0.05	0.0 , 0.02	$6 \times 6 \times 6 \times 6$	0.1~2.5 (for test)
	0.07	0.05 , 0.1	$12 \times 12 \times 12 \times 4$	1.0~2.0(to determine β_c)
	0.1	0.2	$12 \times 12 \times 24 \times 4$	1.1 (for simulations)

Number of configuration for every parameter is also given in Table 2.

Table 2: Statistics of the data

	$\mu = 0.0$	$\mu = 0.02$	$\mu = 0.005$	$\mu = 0.1$	$\mu = 0.2$
$m_q = 0.05$	120	180	160	80	30
$m_q = 0.07$	140	170	160	160	30
$m_q = 0.10$	180	170	160	140	70

Most of the simulations are carried out on small chemical potential and at fixed $\beta = 1.1$ which is little below β_c . Time step of HMC algorithm is $dt=0.01$. After 1000 trajectories of thermalization, data are taken at every 100 trajectories. Since our simulation is restricted in a region of small chemical potential, the algorithm works without difficulty. But, necessary CPU time increases with the chemical potential. A typical trend is shown in table 3. At $\mu = 0.2$, average CPU time is 3.5 times longer than that at $\mu = 0.0$.

Table 3: CPU time of getting one configuration at $\beta = 1.1$, $m_q = 0.05$

	0.0	0.02	0.05	0.1	0.2
time(second)	2223.3	2305.7	2536.0	3122.5	7568.0




Figure 1: (a) $\langle \bar{\psi}\psi \rangle$ vs. β for $L = 6^4$ (diamonds) and $L = 12^3 \times 4$ (circles), (b) Susceptibility of $\langle \bar{\psi}\psi \rangle$ vs. β for $L = 12^3 \times 4$.

4 RESULTS

Firstly we determine phase transition point β_c at $\mu = 0$ on $12^3 \times 4$ lattice.

Chiral condensate $\langle \bar{\psi}\psi \rangle$ versus β is shown in figure 1(a). Chiral restoration occurs at around $\beta \approx 1.4$. In order to determine the critical point more precisely, susceptibility of the condensate is analyzed as shown in figure 1(b). By this analysis, the critical point for $m = 0.07$ is determined as

$$\beta_c = 1.41 \pm 0.03 \quad (5)$$

This value is slightly smaller than that of finite size crossover transition point $\beta_c = 1.54$ on 6^4 lattice.

Nextly, we study quark condensate and pseudoscalar meson mass at small but finite μ . We stay in confinement phase ($\beta = 1.1$). In figure 2(a), we plot the $\langle \bar{\psi}\psi \rangle$ as a function of μ ($0.0 \sim 0.2$) at several m (0.05, 0.07, 0.1). As shown in the figure, the condensate decreases with the chemical potential. But it varies very weakly around $\mu = 0$.

Pseudoscalar meson mass is extracted in a standard way. Typical examples of pseudoscalar correlator is shown in figure 2(b). In the present region of study, single pole fit gives reasonable fitting and results for pion mass squared

are shown in figure 3(a). In order to examine the influence of chemical potential on hadron mass, we examine pseudoscalar meson consisting of light and heavier quarks. figure 3(b) shows their masses at small μ . As shown in the figure, they are very stable and no appreciable dependence on chemical potential is seen in this region.




Figure 2: (a) $\langle \bar{\psi}\psi \rangle$ vs. μ for $m_q = 0.05, 0.07, 0.1$; $\beta = 1.1$. (b) correlation function of $\langle \bar{q}q \rangle$ vs. β for $\beta = 1.1$. $\mu = 0.1$, $m_q = 0.05, 0.07, 0.1$.




Figure 3: (a) M_π^2 vs. m_q for $\beta = 1.1$, $\mu = 0$; (b) M_π vs. μ for $\beta = 1.1$,

5 SUMMARY AND CONCLUSIONS

In this paper, we study 8 flavors SU(2) QCD at finite temperature and small μ on $12^3 \times 4$ and $12^2 \times 24 \times 4$ lattices. The chiral transition at $\mu = 0$ is determined to be $\beta_c = 1.41$. Chiral condensate decreases with μ but variation is very weak around $\mu = 0$. Mass of pseudoscalar meson consisting of light and heavier quarks is stable against chemical potential.

References

1. M. Alford and F. Wilczek, *Phys. Lett. B* **422**, 247 (1999) ;
M. Alford and F. Wilczek, *Nucl. Phys. B* **537**, 443 (1999).
2. R. Rapp, T. Schäfer, E. V. Shuryak, M. Velkovsky, *Phys. Rev. Lett.* **81**, 53 (1998).
3. D. Bailin, and A. Love, *Phys. Rept.* **107**, 325 (1984);
M. Alford, K.Rajagopal, F. Wilczek, *Nucl. Phys. B* **537**, 215 (1999).
4. *see for a review* K. Kanaya, *Nucl. Phys. B* **47**, 47 (1996).
5. A. Nakamura, *Phys. Lett. B* **149**, 391 (1984).
6. S. Duane and J. B. Kogut, *Nucl. Phys. B* **275**, 398 (1986).
7. S. Duane, A. D. Kennedy, B. J. Pendleton, D. Roweth, *Phys. Lett. B* **195**, 216 (1987).
8. C.T.H. Davies, E. G. Klepfish, *Phys. Lett. B* **256**, 68 (1991).
9. S. Gottlieb, W. Liu, R. L. Renken, R. L. Sugar, D. Toussaint *Phys. Rev. D* **38**, 2888 (1988).
10. QCD-TARO Collaboration, *Nucl. Phys. B* **83-84**, 408 (2000).
11. E. B. Gregory, S-H. Guo, H. Kröger, X-Q. Luo, *Phys. Rev. D* **62**, 054508 (2000).
12. S. Hands, J. B. Kogut, M.-P. Lombardo, S. E. Morrison, hep-lat/9902034;
S. Hands, S. E. Morrison, hep-lat/9905021; S. Hands, I. Montvay, S. E. Morrison, M. Pevers, L. Scorzato, J. Skullerud, hep-lat/0006018.
13. I. Montvay, *Nucl. Phys. B* **466**, 259 (1996).
14. T.Hatsuda, S-H.Lee, *Phys. Rev. C* **46**, 34 (1992).